

## What mineral and vitamin levels to recommend in swine diets?

Didier GAUDRÉ, Nathalie QUINIOU

IFIP-Institut du Porc, BP 35104, 35651 Le Rheu Cedex, France  
didier.gaudre@ifip.asso.fr

*Minerals, including trace-elements, and vitamins are currently introduced into pig diets in order to obtain suitable performances. Minerals and vitamins contained in most cereals and usual protein compounds present usually a poor and variable availability. Supplemental diets are then usually needed. This paper, based on literature data, reviews the pig requirement for minerals and vitamins in order to ensure economic and profitable performance. Recent research has focused on phosphorus due to environmental considerations. The widespread use of microbial phytase has decreased the use of inorganic phosphates. Cereals are particularly deficient in calcium and sodium, consequently limestone and salt are necessary to meet the pig requirement for these minerals. Zinc and copper are recognized as growth promoting agents but their use at high levels is detrimental to the environment. The pig diets usually contain supplemental contents of iron, manganese, cobalt, iodine and selenium. The need for these elements in pig diets is described individually. Vitamin E and choline represent the most expensive vitamins added. Choline is then usually not incorporated into the pig fattening diet taking into account the choline contents of the raw materials, while vitamin E is included at a lower level than that which improves the meat quality. Our recent research has dealt with increased levels of vitamins added in combination, compared with most common levels used in farm. Few improvements have been obtained in this way, suggesting that recommendations proposed by INRA in 1984 are mainly justified.*

### Quels niveaux de minéraux et de vitamines recommander pour le régime alimentaire des porcs ?

Les minéraux, les oligo-éléments et les vitamines sont ajoutés couramment à l'alimentation des porcs. La plupart des matières premières disponibles pour l'alimentation des porcs sont caractérisées par des teneurs faibles et des disponibilités réduites et variables en ces éléments. Aussi, des apports complémentaires sont généralement nécessaires sous formes de sels d'oligo-éléments, de vitamines de synthèse et de minéraux. Ce document se base sur une synthèse bibliographique et passe en revue les besoins en minéraux, oligo-éléments et vitamines des porcs permettant d'assurer des performances technico-économiques satisfaisantes dans les conditions d'élevage conventionnelles rencontrées en France. La recherche en nutrition animale a largement orienté ses efforts vers la réduction de l'excrétion de phosphore des porcs, en raison du risque environnemental représenté par son accumulation dans les sols cultivés liés à l'épandage des effluents d'élevage. La généralisation de l'utilisation des phytases microbiennes ces dernières années a entraîné une forte réduction de l'utilisation des phosphates minéraux. Les céréales sont particulièrement déficientes en calcium et en sodium, par conséquent il est nécessaire d'ajouter du carbonate de calcium et du sel pour satisfaire les besoins nutritionnels des porcs. Le zinc et le cuivre sont reconnus comme facteurs de croissance, mais leur utilisation à fortes doses dans les aliments est néfaste pour l'environnement. Les régimes alimentaires des porcs contiennent également des quantités supplémentaires, ajoutées sous forme minérale, de fer, de manganèse, de cobalt, d'iode et de sélénium. Les quantités nécessaires dans les régimes alimentaires des porcs sont décrites séparément pour chaque élément. La vitamine E et la choline sont les vitamines ajoutées les plus onéreuses. Dans ces conditions, il est courant de constater que la choline n'est pas ajoutée aux aliments des porcs à l'engrais, compte tenu des teneurs jugées suffisantes en choline des matières premières. De même, la vitamine E est incorporée à une teneur inférieure à celle permettant d'améliorer la qualité de la viande. Nos travaux récents ont évalué l'intérêt d'apports supplémentaires de vitamines comparé aux pratiques habituelles observées en élevage. Nos résultats montrent peu d'amélioration des performances et nous amènent à considérer que les recommandations proposées par l'INRA en 1984 sont amplement justifiées.

**Keywords:** Feed, mineral, trace-elements, vitamins, performances, requirements

**Mots clés :** Nutrition, minéraux, oligo-éléments, vitamines, performances, besoins

## Introduction

Minerals, including trace-minerals, and vitamins are essential nutrients for pigs. They are usually incorporated into the feed, because common feedstuffs do not have high enough concentration levels in these nutrients, when compared to the needs of the conventional breeding pig. The poor and variable availability of minerals and vitamins from raw materials also support this need. Incorporation rates of added minerals and vitamins represent less than 3% of the feed. Currently, the cost of these ingredients is mainly incurred by phosphates and supplemental vitamins, the prices of which tend to vary greatly especially in recent times.

In practice, vitamins and trace minerals are usually put together into the feed and represent on average, 0.10, 0.15 and 0.30 % of the growing-fattening, sow and piglet feeds, respectively. Macro minerals are introduced separately or could be added to the vitamin and trace mineral pre-mix, in the case of home mixer (25 % of the pig feed production in France). A trend has been observed over recent years, especially in feed manufacture, to decrease the inclusion level of the vitamin and trace element pre-mix (from 1.0 in the early 90's to less than 0.3 % nowadays), in order to reduce the total feed cost. Precautions must be taken to preserve the quality of the feed mixing. This topic must be kept in mind when dealing with mineral and vitamin supplementations.

The aim of the first part of this paper is to review the pig's requirements for macro and trace minerals. Emphasis will be given on minerals with special importance on the environment and/or their cost. The second part deals with pig requirements for vitamins and special attention will be paid on recommended levels to ensure economic performances. It should be pointed out that the purpose of this article is not to describe the metabolic function of each vitamin as this is covered in other research.

## Minerals and trace elements

Pig diets are usually supplemented with limestone, phosphates and salt to meet calcium (Ca), phosphorus (P) and sodium (Na) needs. Table 1 illustrates the deficiencies in Ca and Na of cereals and soybean meal compared to the pig's requirements. Phosphorus is present at higher levels in current raw materials, but at a level of about 60 to 75

percent, in the form of phytate, which is poorly available to the pig. Therefore, inorganic phosphates are added. Since the earlier 90's, microbial phytase has been incorporated into the feed, to improve the availability of P from phytate, through the separation of P from phytate, making it available for intestinal absorption. Recently, the use of such an additive has been generalized and amplified, due to the sharp increase in the price of inorganic phosphates.

**Table 1: Ca, P and Na contents of several raw materials (INRA-AFZ, 2004)**

g/kg	Ca	P	Na
Wheat	0.7	3.2	0.1
Corn	0.4	2.6	0.04
Wheat bran	1.4	9.9	0.1
Soybean meal	3.4	6.2	0.3
Fish meal	38.5	25.2	11.3

Several trace minerals are commonly added to the pig diet to ensure that the pig requirements for iron, copper, manganese, zinc, iodine, cobalt and selenium are met. Raw materials contain insufficient amounts of these nutrients, especially zinc, iodine and selenium (table 2). Furthermore, their availabilities are low and often unknown. Hence, modern pig breeding conditions tend to impose such supplementation in the form of inorganic salts. However, decision makers must be aware that interaction between minerals could lead to a deficiency in one mineral due to the excessive level of another (Perez, 1978): antagonism between copper and iron, copper and zinc, iodine and iron, calcium and copper and calcium and zinc are generally cited.

The content of other minerals i.e. magnesium, potassium, sulphur, chlorine, chromium or molybdenum are much higher in raw materials, compared to the requirements of the pig, specific supplementation is rare.

The environmental risk represented by the accumulation of minerals in the laboured soils has been pointed out over the last few decades, especially in countries where there is a high concentration of animal production. Hence, many studies have focused on the way to reduce the pig excretion of P, copper and zinc. At the same time, national or European regulations have been adopted to limit the dietary incorporation of inorganic sources of these minerals. The trend is not over because mineral content in the plant and exper-

**Table 2: Iron, zinc, manganese, copper, iodine, cobalt and selenium contents of several raw materials (INRA-AFZ, 2004)**

mg/kg	Iron	Zinc	Manganese	Copper	Iodine	Cobalt	Selenium
Wheat	47	27	34	5	0.06	0.02	0.12
Corn	32	19	8	2	0.09	0.05	0.10
Wheat bran	143	74	112	17	0.08	0.09	0.47
Soybean meal	283	47	38	18	0.15	0.26	0.20
Fish meal	351	85	13	7	2.00	0.09	0.40

ted harvest is lower than the level in the liquid manure. Additionally, treatment costs of the slurry are detrimental to the pig industry development.

## Macro Minerals

### Calcium and phosphorus

Calcium and P are essential to the development and constitution of the skeletal system. In the feed, Ca is essentially provided from limestone (on average 38 %). Inorganic phosphates contain high levels of Ca and P (on average 24.5 and 17.5 % of Ca and P for the bi-calcium phosphate). Fishmeal, whey products and meat meal contain high levels of available Ca and P, but their use is limited or even forbidden in western European countries. The Ca and P requirement of the pig is estimated by the factorial method which takes into account the needs for maintenance and production during growth, gestation or lactation periods. According to the factorial approach, the needs during growth are calculated to optimize the skeletal mineralisation (INRA, 1984). A coefficient of net intestinal absorption is then applied to the total need, in order to determine the daily requirement of the pig. This method leads to a Ca requirement of about 7 to 11 g per kg of bodyweight gain, considering a coefficient of net absorption near to 50 % (INRA, 1984). In the case of P, this coefficient is much more variable and depends on the raw materials. Several studies have been conducted over recent decades to estimate P digestibility for the principal components of the pig diet. According to INRA-AFZ (2004), P digestibility coefficients average to 28 and 32 % for corn and soybean meal respectively, which lead to the digestible P contents of 0.7 and 2.0 g/kg. In the case of inorganic sources the digestible coefficients of P are 69 and 83 % for the bi and mono-calcium phosphates respectively. A vegetable phytase is naturally present in straw cereals that theoretically improves the P digestibility from 1.0 to 1.4 g/kg of wheat and 1.1 to 1.4 g/kg of barley. Due to the lack of stability of the vegetable phytase in pellets and insufficient knowledge of its activity and ways to predict its content in raw materials, feed manufacturers usually ignore the contribution of this phytase. Focus is then made on microbial phytases, the activity profile of which has been defined. Table 3 presents those of 3-phytase provided from *Aspergillus Niger* (BASF AG, Ludwigshafen, Germany). It must be highlighted that the P digestible equivalency due to the use of 100 UI/kg feed of phytase, decreases with increasing amounts of phytase. In other words, 1000 UI/kg of phytase are equivalent to 0.95 g/kg of digestible P, when 500 UI/kg represent 0.75 g/kg. Therefore, it is suggested to define several “microbial phytases” in the feedstuffs database used for diet formulation, i.e phytases with decreasing activity and/or digestible P equivalency (Gaudré *et al.*, 2008).

**Table 3: Digestible P equivalencies of the 3-phytase produced by *Aspergillus Niger* from 0 to 1000 FTU/kg feed (Jondreville, 2004)**

g/kg	Digestible P/100 FTU
0 to 250 FTU/kg	0.196
250 to 500 FTU/kg	0.104
500 to 750 FTU/kg	0.052
750 to 1000 FTU/kg	0.028

The digestible P requirement of the growing-fattening pig was estimated by Castaing *et al.* (2003). Based on 17 experiments, they concluded that the optimal growth performance is obtained with 2.5 and 2.0 g/kg feed during the growing (25-65 kg bodyweight) and fattening (65-115 kg) periods respectively. Such levels are in agreement with the estimated values obtained from the factorial approach. Our recent results (Gaudre *et al.*, 2013) indicate that 0.22 and 0.18 g digestible phosphorus per MJ of net energy, could be used respectively in growing and fattening periods. The digestible P requirement is estimated to be 3.5 to 4.0 g/kg for the weaned piglet and the lactating sow, and 2.7 g/kg for the gestating sow.

The metabolisms of Ca and P are closely linked, so it was standard to take into account the Ca/P ratio when formulating the diet. Instead of the Ca/total P ratio, the use of digestible P now involves referring to the Ca/digestible P ratio. According to Jongbloed *et al.* (1999) this ratio should range between 2.7 and 3.2 g/g for piglets and growing pigs. An excess of Ca is particularly detrimental to the phytase activity, making a Ca-phytate combination that is inaccessible to the enzyme, and thus reducing the P digestible equivalency per unit of phytase added.

### Sodium

Pig feeds are currently supplemented with 3 to 5 kg/ton of sodium chloride (38 % Na) to meet the pig's requirement of 1.5 g Na per kg of dry matter feed (INRA, 1984). Despite the fact that the recommendations of Jongbloed *et al.* (1999) are lower (1.0 g/kg feed), it must be emphasised that the Na deficiency led to a loss of appetite. On the other hand, sodium chloride is toxic above 20 kg/ton, especially when water supply is restricted (Perez, 1978) Our results indicate that 3 g of sodium per kg diet with or without 12 g of potassium would not affect pig performances (Alibert, 2012). The amounts of Na, potassium (K) and chlorine (Cl) are usually taken into account together in the dietary electrolyte balance (dEB). The dEB is calculated as  $Na + K - Cl$  and expressed as mEq/kg feed. In this way, the dEB decreases when the diet tends to be acidic. The reduction of total protein contents of the feed leads to a decrease of K amounts. As a consequence, the dEB is reduced. This reduction is even worse when lysine content is kept at the same level through synthetic sources in such diets. Based

**Table 4: Deficiency and sub-deficiency symptoms of trace minerals in the pig (Perez, 1978)**

	Iron	Zinc	Manganese	Copper	Iodine	Selenium
Appetite	*	**			*	
Performances	*	**	*	*	*	*
Fertility disorders		*	**		*	
Skeletal disorders		*	**	**		?
Anaemia	**			*		
Skin or hair anomalies		**		*	*	
Heart disorders				*		
Goitre					**	
Muscular dystrophy						*

on a literature review, Quiniou (2002) indicated that the dEB must be kept between 100 and 275, and 175 and 300 mEq/kg for the weaned piglet and the growing-fattening pigs respectively. Furthermore, a low dEB diet is estimated to induce skeletal demineralisation of the sow (Dourmad & Meschy, 1998). Such a risk sometimes leads to the use of sodium bicarbonate in order to increase the dEB.

### Trace minerals

Under most housing conditions, pigs do not have access to soil or forage. As a consequence, their needs for mineral supplementation may have increased with the generalization of confinement (NRC, 1998). With the exception of iodine (I) for the thyroxin, and iron (Fe) for the haemoglobin, trace minerals are not the constituent of tissue, unlike macro-minerals, but play a catalytic function (Perez, 1978). The role of trace elements in several deficiency symptoms was summarized by Perez (1978, table 4). When estimating the pig's requirements, the occurrence of interactions between trace minerals or between trace minerals and other compounds of the feed must be kept in mind. It should also be noticed that the deficiency symptoms of trace minerals are not specific. For these reasons, recommended levels generally include a safety margin (table 5).

**Table 5: Supplemented trace minerals recommendations (INRA, 1984)**

mg/kg	Piglet	Pig	Sow
Iron	100	80	80
Zinc	100	100	100
Copper	10	10	10
Manganese	40	40	40
Iodine	0.6	0.2	0.6
Cobalt	0.1-0.5	0.1	0.1
Selenium	0.3	0.1	0.1

### Iron

Iron is well known to be the constituent of haemoglobin in the red blood cells, but it is also the component of several molecules, like myoglobin in the muscle or ferritin in the liver. Intra-muscular injection of iron is common practiced to meet the needs of the suckling piglet because the

sow's milk is deficient in this trace mineral. After weaning, the requirement is about 80 mg/kg and tends to decrease during the fattening period in relation to the lower relative content of blood in bodyweight gain (NRC, 1998). Although, contents and availabilities of iron from raw materials are high enough for the pig (NRC, 1998), diets are usually supplied with only 100 mg/kg of iron sulphate in order to avoid any deficiency.

### Zinc

Zinc is the component of many enzymes and thus plays various roles in protein, carbohydrate, lipid and nucleic metabolisms. Many feed constituents are known to increase the need, especially plant phytates and Ca (NRC, 1998). Consequently zinc sulphate or oxide (100 mg/kg) is currently added to pig diets. At pharmacological levels (2000 to 3000 mg/kg), zinc oxide is fed to weaned piglets to prevent diarrhoea and promote growth. Such use is forbidden in France, Germany and the Netherlands while Spain, Denmark, England and Italy have adopted the strategy to temporarily authorize this pharmacological treatment, during the first two weeks after weaning, under veterinarian control. This practice is stigmatised as unfair competition by several feed manufacturers in France, but the adverse effect on the environment must also be pointed out. The retention of zinc by the pig is only 22 mg/kg bodyweight (Dourmad *et al.*, 2002). Hence, almost all of the zinc ingested is excreted in the slurry (Revy *et al.*, 2003). In this context, the maximal zinc content (zinc provided from feedstuffs and mineral supplementation) authorized in the European Union was reduced from 250 to 150 mg/kg in 2003. The use of phytase could also reduce the level of zinc supply. Gaudré *et al.* (2006) demonstrated that 750 UI of 3 phytase produced by *Aspergillus Niger*, can substitute for 40 mg/kg feed of supplemented zinc.

### Manganese

Manganese is essential for the development of the skeleton. It is the component of many enzymes involved in protein, lipid and carbohydrate metabolisms. A manganese deficiency in the diet of the sow could induce reproduction

disorders (Perez, 1978). The NRC's requirement for the sow (20 mg/kg) is based on studies which demonstrated the decrease of piglet birth weight using 5 mg/kg instead of 20 mg/kg, associated to a delayed return to oestrus after weaning (Christianson *et al.*, 1990). Kirchgessner *et al.*, (1981) estimated that the need for manganese is 25 mg/kg for the sow and INRA (1984) recommended 40 mg/kg to avoid any feed deficiency. The usual form of supplementation is manganese oxide.

## Copper

Copper, involved in the defence of the organism against oxidative stress, is required for the synthesis of haemoglobin, and plays a role in the immune system (Jondreville *et al.*, 2002). While the pig's requirement is less than 10 mg/kg feed (NRC, 1998), usual supplementation in piglet diets is currently near 150 mg/kg, due to a growth promoting effect during this stage. The copper retention represents less than 2 mg/kg bodyweight (Jondreville *et al.*, 2002). Therefore, almost all of the copper ingested is then excreted. Regulations on total feed content of copper were adopted in 2003, for environmental reasons. It induced a drastic change on copper supplementation during growing and fattening periods. The maximal authorized content of copper in the feed is now 25 mg per kg of feed, whereas it was 100 mg before 2003. Several veterinarians indicate an increase in gut disorders due to this change, especially at the beginning of the growing phase and in low-clean breeding conditions. It is not known if this regulation induces an increased use of antibiotics. Copper is usually supplied in the form of copper sulphate.

## Iodine

The majority of iodine in swine is present in the thyroid gland as a component of the thyroxine hormone which regulates the metabolic rate (NRC, 1998). The deficiency status led to an increase of piglet stillbirths. The presence of goitrogens substances from several feedstuffs like soybean meal can increase the requirement (NRC, 1998). Cromwell *et al.* (1975) established that the requirement of the growing pig fed a corn soybean meal is approximately 0.09 to 0.14 mg/kg. Safety margins are often used in commercial pig diets through calcium iodate or iodure salts.

## Cobalt

Cobalt is a component of the cyanocobalamin (vitamin B12). There is no evidence that the pig requires cobalt supplements (NRC, 1998). A production activity of cyanocobalamin occurs in the hind gut of the pig (Perez, 1978). Based on cyanocobalamin diet supplementation and pig tendency to coprophagy behaviour, it can be concluded that

diet supplementation has no purpose. For these reasons, sulphate cobalt salt is sometimes omitted from the diet.

## Selenium

As a component of the enzyme glutathione peroxidase, selenium helps cellular membranes to protect against peroxide damage, in association with vitamin E (NRC, 1998). Based on glutathione peroxidase activity in the plasma, Mahan *et al.* (1999) concluded that 0.10 and 0.05 mg/kg of added selenium are sufficient in growing and fattening periods respectively, but they didn't notice any change in performances and carcass characteristics up to 0.3 mg/kg. Compared to a negative control, an improved newly born piglet survival rate is obtained with 0.15 mg/kg of added selenium in the form of sodium selenite or enriched selenium yeast (Mahan & Peters, 2004). No additional effect is observed with 0.3 mg/kg, while the organic source results in greater selenium concentration in the milk. Due to the extremely variable concentration and availability of selenium in soil, several regions of the United States must use selenium supplements to ensure animal health and productivity (Ullrey, 1992). With regard to the risk of selenium deficiency, nutritionists called for regulation in favour of the use of supplemented selenium. It was then authorized in 1974 in the U.S and the maximal authorized level increased from 0.1 to 0.3 mg/kg between 1974 and 1987. The effect of excess dietary selenium on the environment has been pointed out. In the European Union, the maximal content of the feed must not exceed 0.5 mg/kg, while only 5 mg/kg is considered to be toxic for the pig (Perez, 1978).

## Vitamins

Nutrient functions and general considerations on vitamin requirement estimation

Vitamins are essential nutrients for animals. Most of them can be defined as substances that the animal is unable to synthesize, acting in low dose, and not interchangeable (Ferrando *et al.*, 1986). Two groups of vitamins are currently distinguished based on their solubility. Vitamins A, D, E and K are lipid-soluble, whereas B vitamins are water-soluble. Vitamins D and K are involved in mineral metabolism and blood coagulation respectively. Vitamins A and E are involved in a lot of metabolism pathways, and in consequence could be implicated in overall performances. Group-B vitamins are co-enzymes required in numerous metabolisms (carbohydrates, lipids, proteins, nucleic acids).

Feedstuffs contain vitamins, but it must be underlined that their contributions are very difficult to take into account. The amount of vitamins in feedstuffs varies between publications. Pyridoxine (B6) exists in the form of three active

compounds that complicate laboratory analyses (INRA, 1984). It could also vary between crops, harvest and variety, as it is the case for biotin (H) from cereals (INRA, 1984). Vitamins from raw materials could exist in different molecular forms with varied activities. The L-isomer of biotin is a biologically inactive form for the pig (NRC, 1998). Vitamin E is present in feedstuffs in eight natural forms ( $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  tocopherols and tocotrienols), each characterized by its own vitamin E activity. Interactions with feed components also need to be mentioned, because they can increase or decrease the need for supplementation. The presence of methionine reduces the need for cyanocobalamin (B12), as does the presence of tryptophane for niacin (PP). Niacin in cereals is largely unavailable to young pigs (NRC, 1998). Furthermore, it must be specified that the vitamins naturally present in raw materials are unstable, especially in the case of tocopherols and tocotrienols, and also with thiamin (B1) when heat treatment is applied.

The gastro intestinal tract is colonized by numerous bacterial populations. A production activity of vitamins K, folic acid (B9), cyanocobalamin, niacin, and biotin can be observed in the large intestine. Except for vitamin K which can be directly digested, the absorption of B vitamins needs coprophagy. The generalization of slatted floors in modern breeding conditions, tends to reduce this vitamin supply.

However, the contribution from raw materials and intestinal syntheses is usually considered in the case of thiamin, riboflavin (B2), pantothenic acid (B5), niacin, biotin, choline, pyridoxine, folic acid and vitamin E provided from raw materials, and vitamins K, folic acid, cyanocobalamin, biotin and niacin due to intestinal bacterial activity.

Investigations into vitamin needs of animals are particularly complex. Deficiency symptoms are not specific, especially for water-soluble vitamins. Synthetic diets are necessary to avoid the presence of vitamins from raw materials, and attention must be paid on interaction with intestinal synthesis, and with other nutrients that increase or decrease the need. Liver storage would interact with requirement assessments for vitamins A, E and B12, by buffer effect. The variability of needs among individuals and evolution of needs with genetic improvement must also be emphasized; most of the determination of vitamin requirements were conducted in the 40's and 50's, with an animal model far removed from our current pig. The observed performances are considered to be, in several cases, insufficient criteria to estimate the real need. Other parameters provided from blood, bone, liver, colostrum or milk are therefore measured to complete the data. Such parameters are chosen in relation to the metabolic role of the vitamin studied. Sometimes, the levels optimising performances are lower than those optimising the parameter. In such a case, deci-

sion makers must be aware of the supplemental cost of the vitamin.

## Literature review

This literature review aims to point out noteworthy results that could help the reader to define diet supplemental levels of vitamins needed by the modern breeding herd.

Cure treatment of vitamins is usually practiced on the sow herd, but apparently it seems unjustified because, except for vitamin B12, liver storage of vitamin B doesn't exist. Additionally, liver storage of vitamin E is limited. Then most of the studies and recommendations focus on the daily rate and consequently, regular ingestion of vitamins.

### Vitamin A

Vitamin A supplementations of 40000 UI/kg for piglets and 50000 UI/kg for pigs failed to improve feed efficiency or average daily gain. On the other hand, small amounts (<500 UI/kg) are enough to optimise performances. Long term studies on sows are complicated by the high storage capacity of the liver. Due to the oxidation sensitivity of natural and in a lesser extent commercial sources of vitamin A, supplementation is usually kept at a high level, knowing that the cost of 5000 UI/kg is not expensive (approximately 0.17 €/T).

### Vitamins D and K

The pig's ability to cope with a deficiency of vitamin D is high (Pointillart, 1980), while intestinal synthesis of vitamin K is able to meet the body's requirements. However, because the supplementation of vitamins D and K is cheap, levels are usually near 2000 UI/kg for vitamin D and between 0.5 to 1.0 mg/kg for vitamin K.

### Vitamin E

Vitamin E is the most expensive vitamin. Every 10 UI/kg costs approximately 0.35 €/ton. The need for vitamin E for the growing pig was reviewed by Ullrey (1981) on the basis of 22 publications. He recommended a level of vitamin E between 10 and 20 UI per kg feed and even 30 UI/kg when a problem occurred in the breed. At the opposite end of the scale, 5 UI/kg appeared insufficient. Conclusions drawn by Jensen *et al.* (1988) were similar; 15 UI/kg are suitable, 30 UI/kg could be used taking into account the variability of the need between pigs and including a safety margin. The effect of vitamin E on meat quality has been the subject of many studies. Unfortunately, an expensive level of vitamin E (100 to 200 UI/kg) is needed to modify the colour of meat and/or resistance to oxidation. Antagonism between vitamins A and E was demonstrated by Ching *et al.* (2002) on piglets. Blood content of tocopherol was reduced from 20 to 30 % when the A supplementation increased from 2200 to 13200 UI/kg. This antagonism must be kept in mind even

if it hasn't been confirmed on fattening pigs with 20000 UI/kg of vitamin A (Anderson *et al.*, 1995). Vitamin E supplementation for sows has increased in France since the early 90's (approximately, 20 to 80 UI/kg), in response to the widespread use of hyper-prolific sows. Also, the simultaneous reduction of its cost might help this trend. In fact, demonstration of such a high supplementation is scarce. Mahan (1994) recommended 44 UI/kg for both gestating and lactating sows based on a slight increase of the number of piglets born alive.

### Thiamin

Thiamin is abundant in cereals and soybean meal (> 4 mg/kg). Whatever the physiological stage, experiments failed to improve performances, when more than 1 mg/kg of supplemented thiamin was added (Newcomb & Allee, 1986; Woodworth *et al.*, 2000, Easter *et al.*, 1983; Lutz *et al.*, 1999).

### Riboflavin

Recent works have been conducted on riboflavin. Lutz & Stahly (1998) showed a linear increase of average daily gain, feed efficiency and body protein accretion of high-lean piglets with 3.7 and 7.4 mg/kg of riboflavin added (table 6). They determined that the riboflavin need, necessary for the protein accretion, is six times higher than that necessary for the fat deposit. In other words, riboflavin need is directly dependent on the genetic potential of the pigs. Unfortunately, no significant changes in pig fattening performances were observed (Gaudré & Granier, 2009) with added riboflavin up to 18 mg/kg (table 7). But it must be pointed out that in this experiment, the lowest level was 3 mg/kg of riboflavin added, plus 1.6 mg/kg provided from feedstuffs.

**Table 6: Effect of added riboflavin on piglet performance Lutz & Stahly (1998)**

Added riboflavin mg/kg	0	3.7	7.4
ADG (g/d)	645	677	696
FCR (kg/kg)	1.43	1.36	1.34
Protein accretion (g/d)	110	115	118

**Table 7: Effect of added riboflavin on pig performance Gaudré & Granier (2009)**

Added riboflavin mg/kg	3	6	9	18
ADG (g/d)	938	961	946	927
FCR (kg/kg)	2.80	2.80	2.80	2.74
Lean thickness (mm)	56.0	57.4	56.7	57.4

### Pantothenic acid

Slight improvements in performance are obtained with pantothenic acid addition. Grinstead *et al.* (1988) showed on piglets weaned at 11 days old, an increase in average daily gain with 120 mg/kg of pantothenic acid added and in feed efficiency with 30 mg/kg. Stahly & Lutz (2001) impro-

ved lean percent carcass with 30 mg/kg without any other change in fattening performances observed.

### Pyridoxine

Woodworth *et al.* (2000) improved the average daily gain of weaned piglets from day 0 to 14, with 3.3 mg of pyridoxine added and 4.6 mg provided from feedstuffs. They concluded that the requirement of the weaned piglet is nearer to 7.9 mg/kg. Unfortunately, this positive effect was not observed by Newcomb & Allee (1986) with the same supplemental level.

### Folic acid

Because of the contribution by raw materials and intestinal synthesis, the need for folic acid is considered as low, with the exception of the young animal and the pregnant sow. The addition of folic acids on piglets and pigs were tested, without any positive change, by Newcomb & Allee (1986), Easter *et al.* (1983) and Lutz *et al.* (1999). Sow prolificacy is not always increased by acid folic additions: Harper *et al.* (1994) with 1, 2 or 4 mg/kg, Matte *et al.* (1992) with 5 or 15 mg/kg throughout pregnancy. This effect seems dependent on the ovulation rate (Lindemann & Kornegay, 1989), since the latter increases with parity.

### Cyanocobalamin

Simard *et al.* (2004) observed improvements in total piglets born and weaned with high levels of vitamin B12 (cyanocobalamin) in the diets of pregnant sows (0.2 and 0.4 mg/kg). The piglet hepatic storage of B12 tends to increase at the same time. It should be highlighted that this test was carried out with only 8 sows per treatment, and needs to be confirmed on a larger scale. Furthermore, the cost of 0.2 to 0.4 mg/kg of cyanocobalamin is similar to the addition of 12 to 24 UI/kg of vitamin E. The weaned piglet didn't show any increase in its performance with up to ten times the NRC requirement (Lindeman *et al.*, 1995).

### Niacin

Niacin supplementation is required when the content of niacin calculated from raw materials, is lower than 1.5 times the minimal need (INRA, 1984): 33, 18 and 15 mg/kg for piglet, pig and sow, respectively. This recommendation is based on the low availability of niacin from raw materials (corn especially), and the presence of natural antagonism from niacin. No effect was observed by Copelin *et al.* (1980) and Ivers & Veum (1993) up to 22 and 81 mg/kg of added niacin respectively, while Real *et al.* (2002) demonstrated improved growth and feed efficiency performance with 13 to 55 mg/kg.

### Biotin

The interest of biotin supplementation for weaned piglets and growing-finishing pigs was not demonstrated

**Table 8: Effects of biotin supplementation on sow productivity**

Authors	mg/kg	Noticed improvement
Hamilton & Veum (1984)	0.55	Piglets weaned on first parity
Bryant et al. (1985b,c)	0.44	Trend on prolificacy, weaning-oestrus interval and foot lesions
Lewis et al. (1991)	0.33	Weaned/born alive
Watkins et al. (1991)	0.44	Nothing

(Newcomb & Allee, 1986; Easter *et al.*, 1983; Hamilton & Veum, 1986; Bryant *et al.*, 1985a). On the other hand, several studies tend to conclude that biotin supplementation needs to be added to the feed for the sow (Hamilton & Veum, 1984; Bryant *et al.*, 1985b,c; Lewis *et al.*, 1991) with the exception of Watkins *et al.* (1991). Results indicate that 0.3 to 0.55 mg/kg could enhance sow performance (table 8).

### Choline

Choline is not strictly considered as a vitamin but supplementation could be efficient to complement choline provided from raw materials. The sow requirement is 2000 mg/kg taking into account each source of choline (INRA, 1984). Additionally, NRC (1998) indicates that supplemental choline is not necessary for growing-fattening pigs eating a diet rich in soybean meal. Choline is very expensive (300 mg/kg induce 0.5 €/T) and is usually restricted to the sow and piglet feed.

### Effects of vitamins added in combination

During the last few years, IFIP has carried out several studies on vitamin requirements. The approach was to estimate the effect of vitamins added in combination. In these experiments, the control treatment was considered to be represented by usual practice. The growing-fattening trial (Gaudré & Vautier, 2006) aimed at comparing a high level of combined vitamins (table 9), but no significant improvement was observed on breeding performances and carcass characteristics. A tendency to enhance average

daily gain and feed intake of piglets weaned in low-clean conditions, was demonstrated due to a moderate increase in added vitamins (table 10). This result is similar to those published recently with added vitamins B (Wilson *et al.*, 1992a,b; Lindemann *et al.*, 1995; Stalhy *et al.*, 1995) or B and fat-soluble vitamins (Castaing *et al.*, 2001). In a cooperative study, involving 336 sows from two herds (Quiniou & Calvar, 2005), no improvement in productivity was noticed when using higher than recommended levels of vitamin E, K and a range of B vitamins (table 11). Before publication of these works, there was a trend in Brittany to increase feed vitamin supplementation, believing that common practice of feed manufacturers led to nutrient deficiencies in pigs. It was demonstrated that little improvement in performances could be obtained in this way.

### Levels of vitamins recommended by phase

Based on review literature and our own recent investigations described above, recommendations were formulated for added vitamins (table 12). These recommendations are partly based on those established by INRA in 1984. Bearing in mind the various sources of incertitude, intervals of variation are sometimes indicated. In some cases, recommended levels are made but they are based on a limited number of trials. Taking into considerations, the price and time-varying of vitamins, feed components, the use of safety margins, as well as newly acquired knowledge, judgements could be arrived at in favour of contribution levels different from those suggested here.

**Table 9: Supplemented vitamin levels compared on growing-fattening pigs (Gaudré & Vautier, 2006)**

Vitamins	Unit/kg	Growing period (25-60 kg)		Fattening period (60-115 kg)	
		Control	Test	Control	Test
A	UI	6260	10000	5700	8000
D3	UI	1200	2000	1075	1500
E	UI	9	60	8	50
K3	mg	0.26	3	0.12	1.5
B1 (Thiamin)	mg	0.5	2	0.4	1.5
B2 (Riboflavin)	mg	2.3	10	2	8
B5 (Pantothenic Ac.)	mg	7	40	6	40
B6 (Pyridoxine)	mg	0.5	4	0.4	3
B12 (Cyanocobalamin)	mg	0.014	0.04	0.012	0.035
PP (Niacin)	mg	11	30	10	30
Folic Ac.	mg	0	1	0	1
Biotin	mg	0	0.25	0	0.2
Choline	mg	300	300	200	200

**Table 10: Supplemented vitamin levels compared on piglets (Gaudré, 2007)**

Vitamins	Unit/kg	Control	Test
A	UI	8000	12000
D3	UI	1500	2000
E	UI	20	60
K3	mg	0.5	2
B1 (Thiamin)	mg	1	2
B2 (Riboflavin)	mg	4	8
B5 (Pantothenic Ac.)	mg	10	20
B6 (Pyridoxine)	mg	1.5	3
B12 (Cyanocobalamin)	mg	0.015	0.03
PP (Niacin)	mg	15	45
Folic Ac.	mg	0.5	1
Biotin	mg	0	0.1
Choline	mg	300	450

**Table 11: Supplemented vitamin levels compared on sows (Quiniou & Calvar, 2005)**

Vitamins	Unit/kg	Control	Test
A	UI	9600	9600
D3	UI	1960	1960
E	UI	50	90
K3	mg	0.5	1
B1 (Thiamin)	mg	0.6	1.5
B2 (Riboflavin)	mg	4.2	6
B5 (Pantothenic Ac.)	mg	10	20
B6 (Pyridoxine)	mg	0.6	2.5
B12 (Cyanocobalamin)	mg	0.025	0.025
PP (Niacin)	mg	17	35
Folic Ac.	mg	1.8	3.5
Biotin	mg	0.15	0.30
Choline	mg	600	600

**Table 12: Recommended levels for vitamins (IFIP, 2005)**

Vitamins Units/kg	Cost	Piglet		Pig		Sow	
		7-12 kg BW	12-25 kg BW	25-60 kg BW	60-115 kg BW	Pregnant	Lactating
A UI	\$\$	10000	8000	7000	5000	5000	8000
D3 UI	\$	2000	1500	1000	1000	2000	2000
E UI	\$\$\$	50	20	10-15	10-15	45	45
K3 mg	\$	1	0.5-1	0.5	0.5	0.5	0.5
B1 (Thiamin) mg	\$	1	0.5-1	0-1	0-1	1	1
B2 (Riboflavin) mg	\$\$	4-8*	4-8*	3	2	3	3
B5 (Panto. ac.) mg	\$\$	10	10	8	7	8	8
PP (Niacine) mg	\$\$	15	15	10	7	10	10
B6 (Pyridoxine) mg	\$\$	0-3.6*	0-1	0	0	1	0
B12 (Cyanocobalamin) mg	\$\$	0.035	0.03	0.02	0.02	0.02-0.2	0.02
Folic ac. mg	\$\$	0.5	0.5	0-0.5	0-0.3	0.2-15*	1-15*
Biotin mg	\$\$\$	0.1	0-0.1	0-0.05	0	0.3-0.5	0.3-0.5
Choline mg	\$\$\$	300-800	300-800	0-300	0-100	400-900	400-900

\*: these levels are based on a limited number of trials

## Conclusion

The inclusion of minerals, including trace elements, and vitamins in pig diets is needed to ensure optimal performance of the actual pig genotype deprived of access to soil or forage. Recent knowledge has been collected on phosphorus, zinc and copper in response to increasing awareness of detrimental effects of the excess of these minerals on the environment. As a consequence, investigations have been conducted in order to limit the total excretion of the pig, considering that the retention of phosphorus, zinc and copper is generally not affected by diet content. Partly due to a decrease in the protein in the diet and thus potassium content, attention has been paid on the dietary electrolyte balance taking into account respective levels of

sodium, potassium and chlorine, but real proof of their reliability remains scarce.

Over the last two decades, focus has been made on vitamin requirements, since most of the research conducted in the 40's and 50's, was based on an animal model far removed from our current pig. Several studies have been carried out in order to estimate if improved performance could be obtained at higher vitamin levels than current recommendations. Our recent research and literature review have been summarized in practical recommendations presented above. Safety margins could be introduced bearing in mind the actual cost of each vitamin.

## References

- Alibert, L. Influence de l'excès de sodium et potassium sur la consommation et la croissance des porcs charcutiers. *Tech porc*, N° 3, P. 12-13, 2012.
- Anderson, L.E.; Myer, R.O.; Brendemuhl, J.H.; Mc Dowell L.R. The effect of excessive dietary vitamin A on performance and vitamin E status in swine diets varying in dietary vitamin E. *Journal of Animal Science*, v.73, p.1093-1098, 1995.
- Bryant, K.L.; Kornegay, E.T.; Knight, J.W.; Webb, K.E.; Notter, D.R. Supplemental biotin for swine. I. Influence on feedlot performance, plasma biotin, and toe lesions in developing gilts. *Journal of Animal Science*, v.60, n.1, p.136-144, 1985a.
- Bryant, K.L.; Kornegay, E.T.; Knight, J.W.; Webb, K.E.; Notter, D.R. Supplemental biotin for swine. II. Influence of supplementation to corn and wheat based diets on reproductive performance and various biochemical criteria of sows during four parities. *Journal of Animal Science*, v.60, n.1, p.145-153, 1985b.
- Bryant, K.L.; Kornegay, E.T.; Knight, J.W.; Webb, K.E.; Notter, D.R. Supplemental biotin for swine. III. Influence of supplementation to corn and wheat based diets on the incidence and severity of toe lesions, hair and skin characteristics and structural soundness of sows housed in confinement during four parities. *Journal of Animal Science*, v.60, n.1, p.154-162, 1985c.
- Castaing, J.; Cambelil, D.; Hamelin, C. Incidence des apports en vitamines sur les performances des porcelets en deuxième âge. *Journées de la Recherche Porcine*, v.33, p.227-232, 2001.
- Castaing, J.; Paboeuf, F.; Skiba, F.; Chauvel, J.; Cazaux, J.G.; Milgen, J.; Jondreville, C. Estimation du besoin en phosphore digestible apparent du porc charcutier : synthèse d'essais zootechniques effectués au cours des dix dernières années. *Journées de la Recherche Porcine*, v.35, p.47-54, 2003.
- Ching, S.; Mahan, D.C.; Wiseman T.G.; Fastinger, N.D. Evaluating the antioxidant status of weanling pigs fed dietary vitamins A and E. *Journal of Animal Science*, v.80, p.2396-2401, 2002.
- Christianson, S.L.; Peo, E.R.; Lewis, A.J.; Giesemann, M.A. Influence of dietary manganese levels on reproduction, serum cholesterol and milk manganese concentration of sows. *Journal of Animal Science*, v.68, Suppl.1, Abstracts, p.368, 1990.
- Copelin, J.L.; Monegue, H.J.; Combs, G.E. Niacine levels in growing-finishing swine diets. *Journal of Animal Science*, abstracts, p.190, 1980.
- Cromwell, G.L.; Sihombing, D.T.H.; Hays, V.W. Effects of iodine level on performance and thyroid traits of growing pigs. *Journal of Animal Science*, v.41, n.3, p.813-818, 1975.
- Dourmad, J.-Y.; Meschy, F. Le bicarbonate de sodium en nutrition porcine. In : Conférence GTV/SPACE, Rennes, France, 1998.
- Dourmad, J.-Y.; Pomar, C.; Masse, D. Modélisation du flux de composés à risque pour l'environnement dans un élevage porcin. *Journées de la Recherche Porcine*, v.34, p.183-194, 2002.
- Easter, R.A.; Anderson, P.A.; Michel, E.J.; Corley, J.R. Response of gestating gilts and starter, grower and finisher swine to biotin, pyridoxine, folacin and thiamine additions to corn-soybean meal diets. *Journal of Animal Science*, v.28, n.5, p.945-954, 1983.
- Ferrando, R.; Ruckebusch, Y.; Rerat, A. Les vitamines. Le porc et son élevage, bases scientifiques et techniques. Ed. Maloine. 1986. 575p.
- Gaudre, D.; Vautier, A. Incidence zootechnique d'un taux de complémentation vitaminique élevé en engraissement. *Techniporc*, v.29, n.2, p.19-26, 2006.
- Gaudre, D.; Larrere, V.; Granier, R.; Magnin, M.; Jondreville, C.; Paboeuf, F. Quelle réduction du rejet de zinc la 3-phytase microbienne permet-elle chez le porc à partir de 12 kg de poids vif ? *Journées de la Recherche Porcine*, v.38, p.5-12, 2006.
- Gaudre, D. Incidence zootechnique d'une augmentation modérée de la complémentation vitaminique de l'aliment distribué entre 12 et 30 kg de poids vif. *Techniporc*, v.30, n.1, p.21-28, 2007.
- Gaudre, D.; Alibert, L.; Quiniou, N.; Royer, E. Recommandations et calcul de l'apport en phosphore digestible par les matières premières et les phytases. *Techniporc*, v.31, n.5, p.33-39, 2008.
- Gaudre, D.; Granier, R. Incidence de la riboflavine sur les performances des porcs en engraissement. *Journées de la Recherche Porcine*, v.41, p.139-140, 2009.
- Gaudré, D., Monziols, M., Lebas, N. Incidence du rapport phosphore : énergie de l'aliment sur les caractéristiques du radius mesurées au scanner à rayons X. *Journées de la recherche porcine*, P. 199-200, 2013.
- Grinstead, G.S.; Goodband, R.D.; Nelssen, J.L.; Tokach, M.D., Dritz, S.S.; Stott, R. Effects of increasing pantothenic acid on growth performance of segregated early-weaned pigs. *Swine Day 1998*, p.88-90.
- Hamilton, C.R.; Veum, T.L. Responses of sows and litters to added dietary biotin in environmentally regulated facilities. *Journal of Animal Science*, v.59, n.1, p.151-157, 1984.
- Hamilton, C.R.; Veum, T.L. Effect of biotin and (or) lysine additions to corn-soybean meal diets on the performance and nutrient balance of growing pigs. *Journal of Animal Science*, v.62, p.155-162, 1986.
- Harper, A.F.; Lindemann, M.D.; Chiba, L.I.; Combs, G.E.; Handlin, D.L.; Kornegay, E.T.; Southern, L.L. An assessment of dietary folic acid levels during gestation and lactation on reproductive and lactational performance of sows: a cooperative study. *Journal of Animal Science*, v.72, p.2338-2344, 1994.
- IFIP-French Institute for pig and pork industry. Vitamines : rôles et besoins dans l'aliment du porc. Ed. IFIP, Paris, 2005.
- INRA. L'alimentation des animaux monogastriques: porc, lapin, volailles. Ed. INRA, Paris, 1984.
- INRA-AFZ. Tables de composition et de valeur nutritive des matières premières destinées aux animaux d'élevage. Ed. INRA, Paris, 2004.
- Ivers, D.J.; Veum, T.L. Effect of niacin additions to corn-soybean meal diets on performance of pigs from weaning to finishing. *Journal of Animal Science*, v.71, p.3383-3388, 1993.
- Jensen, M.; Hakkarainen, J.; Lindholm, A.; Jönsson, L. Vitamin E requirement of growing swine. *Journal of Animal Science*, v.66, p.3101-3111, 1988.
- Jondreville, C.; Prevy, P.S.; Jaffrezic, A.; Dourmad, J.-Y. Le cuivre dans l'alimentation du porc : oligo-élément essentiel, facteur de croissance et risque potentiel pour l'homme et l'environnement. *INRA Productions Animales*, v.15, n.4, p.247-265, 2002.
- Jondreville, C. Calculer la teneur en P digestible apparent d'un aliment pour porcs à partir des tables INRA-AFZ, Addendum, 2004.
- Jongbloed, A.W.; Everts, H.; Kemme, P.A.; Mroz, Z. Quantification of absorbability and requirements of macroelements In: KYRIAZAKIS, I. (Ed.) A quantitative biology of the pig, CAB International, 1999, p.275-298.
- Kirchgessner, M.; Roth-Maier, D.A.; Sporl, R. Untersuchungen zum Trachtigkeitsanabolismus der spurenelemente Kupfer, Zin, Nickel und Mangan bei Zuchtsaver. *Arch.Tierernährung*, v.31, p.21-34, 1981.
- Lewis, A.J.; Cromwell, G.L.; Pettigrew, J.E. Effects of supplemental biotin during gestation and lactation on reproductive performance of sows: a cooperative study. *Journal of Animal Science*, v.69, p.207-214, 1991.
- Lindemann, M.D.; Kornegay, E.T. Folic acid supplementation to diets of gestating-lactating swine over multiple parities. *Journal of Animal Science*, v.67, p.459-464, 1989.

- Lindemann, M.D.; Cromwell, G.L.; Monegue, H.J. Effects of inadequate and high levels of vitamin fortification on performance of weanling pigs. *Journal of Animal Science*, v.73, suppl.1, p.16, 1995.
- Lutz, T.R.; Stahly T.S. Dietary riboflavin needs for body maintenance and body protein and fat accretion in pigs. *Iowa State University*, p.41-44, 1998.
- Lutz, T.R.; Stahly T.S.; Cook, D.R.; Ewan, R.C. Effects of dietary thiamin, folacin or niacin regimen on growth in high lean pigs. *Journal of Animal Science*, v.77, suppl.1, p.189, 1999.
- Mahan, D.C. Effects of dietary vitamin E on sow reproductive performance over a five-parity period. *Journal of Animal Science*, v.72, p.2870-2879, 1994.
- Mahan, D.C.; Cline, T.R.; Richert, B. Effects of dietary levels of selenium-enriched yeast and sodium selenite as selenium sources fed to growing-finishing pigs on performance, tissue selenium, serum glutathione peroxidase activity, carcass characteristics and loin quality. *Journal of Animal Science*, v.77, p.2172-2179, 1999.
- Mahan, D.C.; Peters, J.C. Lon-term effects of dietary organic and inorganic selenium sources and levels on reproducing sows and their progeny. *Journal of Animal Science*, v.82, p.1343-1358, 2004.
- Matte, J.J.; Girard, C.L.; Brisson, G.J. The role of folic acid in the nutrition of gestating and lactating primiparous sows. *Livestock Production Science*, v.32, p.131-148, 1992.
- Newcomb, M.D.; Allee, G.L. Water soluble vitamins for weaned pigs. *Journal of Animal Science*, v.64, suppl.1, p.108, 1986.
- NRC. Nutrient Requirement of Swine, tenth revised edition, National Academy Press, Washington, D.C., 1998.
- Perez, J.M. L'alimentation minérale des porcins. *Techniporc*, v.1, n.1, p.1-32, 1978.
- Pointillart, A. La carence expérimentale en vitamine "D" chez le porc. *Journées de la Recherche Porcine en France*, v.12, p.335-344, 1980.
- Quiniou, N. Le point sur l'équilibre acido-basique chez le porc et le bilan électrolytique des aliments. *Techniporc*, v.25, n.3, 2002
- Quiniou, N.; Calvar, C. Est-ce que la truie hyperprolififère valorise un apport en vitamines supérieur aux recommandations ? *Techniporc*, v.28, n.5, p.3-8, 2005.
- REAL, D.E.; Nelssen, J.L.; Unruh, J.A.; Tokach, M.D.; Goodband, R.D.; Dritz, S.S.; Derouchey, J.M.; alonso, E. Effects of increasing dietary niacin on growth performance and meat quality in finishing pigs reared in two different environments. *Journal of Animal Science*, v.80, p.3203-3210, 2002.
- Revy, P.S.; Jondreville, C.; Dourmad, J.-Y.; NYS, Y. Le zinc dans l'alimentation du porc : oligo-élément essentiel et risque potentiel pour l'environnement. *INRA Productions Animales*, v.16, n.1, p.3-18, 2003.
- Simard, F.; Guay, F.; Girard, C.L.; Giguere, A.; Laforest, J.-P.; Matte, J. La vitamine B12 chez la truie gravide : faut-il en actualiser le besoin ? *Journées de la Recherche Porcine*, v.36, p.229-234, 2004.
- Stahly, T.S.; Williams, N.H.; Swenson, S.G.; Ewan, R.C. Dietary B vitamin needs of high and moderate lean growth pigs fed from 9 to 28 kg body weight. *Journal of Animal Science*, v.73, suppl.1, p.193, 1995.
- Stahly, T.S.; Lutz, T.R. Role of pantothenic acid as a modifier of body composition pigs. *Journal of Animal Science*, v.79, suppl.1, p.68, 2001.
- Ullrey, D.E. Vitamin E for swine. *Journal of Animal Science*, v.53, n.4, p.1039-1056, 1981.
- Ullrey, D.E. Basis for regulation of selenium supplements in animal diets. *Journal of Animal Science*, v.70, p.3922-3927, 1992.
- Watkins, K.L.; Southern, L.L.; Miller, J.E. Effect of dietary biotin supplementation on sow reproductive performance and soundness and pig growth and mortality. *Journal of Animal Science*, v.69, p.201-206, 1991.
- Wilson, M.E.; Pettigrew, J.E.; Johnston, L.J.; Chester-Jones, H. Effect of B vitamin supply upon growth of weanling pigs. *Journal of Animal Science*, v.70, suppl.1, p.61, 1992a.
- Wilson, M.E.; Pettigrew, J.E.; Shurson, G.C.; Johnston, L.J.; Chester-Jones, H. Effect of additional B vitamin in complex diets containing porcine plasma proteins and milk products or in simple diets on growth of weanling pigs. *Journal of Animal Science*, v.70, suppl.1, p.233, 1992b.
- Woodworth, J.C.; Goodband, R.D.; Nelssen, J.L.; Tokach, M.D.; Musser, R.E. Added dietary pyridoxine, but not thiamine, improves weanling pig growth performance. *Journal of Animal Science*, v.78, p.88-93, 2000.

## Reference to this article

- Gaudré D., Quiniou N., 2015. What mineral and vitamin levels to recommend in swine diets?. *Les Cahiers de l'IFIP*, 2(1), 49-59.

